




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : A61F 13/00, A61L 15/01	A1	(11) International Publication Number: WO 87/05206 (43) International Publication Date: 11 September 1987 (11.09.87)
<p>(21) International Application Number: PCT/SE87/00113</p> <p>(22) International Filing Date: 6 March 1987 (06.03.87)</p> <p>(31) Priority Application Number: 8601098-0</p> <p>(32) Priority Date: 10 March 1986 (10.03.86)</p> <p>(33) Priority Country: SE</p> <p>(71) Applicant (for all designated States except US): MÖLN- LYCKE AB [SE/SE]; S-405 03 Göteborg (SE).</p> <p>(72) Inventor; and</p> <p>(75) Inventor/Applicant (for US only) : FABO, Tomas [SE/ SE]; Gråbergsvägen 13, S-435 00 Mölnlycke (SE).</p> <p>(74) Agents: HJÄRNE, Per-Urban et al.; H. Albihns Patent- byrå AB, Box 7664, S-103 94 Stockholm (SE).</p>		<p>(81) Designated States: AT (European patent), BE (Euro- pean patent), CH (European patent), DE (European patent), DK, FI, FR (European patent), GB (Euro- pean patent), IT (European patent), JP, LU (Euro- pean patent), NL (European patent), NO, SE (Euro- pean patent), US.</p> <p>Published <i>With international search report.</i></p>

(54) Title: WOUND DRESSING**(57) Abstract**

Soft, elastic and flexible wound dressing having through-holes (5) or channels admitting the passage of fluid and air therethrough. The dressing is composed of a hydrophobic gel (2) applied to a reinforcement (1) made of an elastic material.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
RJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

WOUND DRESSING

A wound healing process may principally be subdivided into three phases. The wound is first cleaned out followed by the formation of new tissue, whereafter the restored tissues during a final maturing phase stabilize while developing a less brittle and more elastic structure.

The regeneration phase involves the growth of capillaries, fibroblasts and epithelium into the wound site for building up new tissue. The newly formed tissue is extremely delicate and supersensitive to external influences. If a wound still in progress of regenerating tissue is covered with a dressing composed of a fibrous material, the fibers may easily intermingle with the newly formed tissues and give rise to inflammatory reactions in the wound tissue, which would result in deterioration of the wound healing process. Furthermore, the wound tissue would also be mechanically damaged in connection with removal and change of dressing. To avoid this, it is extremely important for the dressing applied to the wound not to get stuck in dried-up wound exudate, or in any coagulum possibly formed. To attain the most favorable conditions for optimal wound healing, the wound should be kept moist but free of excess wound exudate during this phase of wound repair.

Wound dressings intended for use during this particularly sensitive stage of the wound healing process should therefore be designed so as not to stick to the wound bed; they should be pliable and have a soft wound-contacting surface. In addition, the dressings must be capable of seeping up excess amounts of wound fluid, or to allow for the passage of fluid exudate into an absorbent body placed over the dressing.

Commonly used types of dressings for application to wounds in the regenerating phase are pads soaked with ointment and made of gauze or nylon netting, possibly in

combination with an absorbent body. Dressings of this type, as compared with conventional fibrous dressings, have a low tendency of sticking to the wound due to their poor adhesivity towards wound tissue. However, there are associated with the use of such dressings numerous drawbacks and disadvantages. For example, ointment easily falls off during use of this dressing admitting thereby a foreign substance to enter the wound, which would negatively affect the wound healing process. The pads soaked with ointment are smeary and unpleasant to handle and, although impregnated, they often stick tenaciously to the wound bed thus giving rise to tissue damages.

Another type of dressing used on wounds in the regenerating stage constitutes a combination of a perforated polymer or metal film and a more or less absorbent body made of a fibrous material. The idea is for the film to produce a non-frictional, hydrophobic wound-contacting surface preventing in this manner the dressing from sticking to the wound. Dressings of this type are stiff, inelastic and non-flexible. The greatest disadvantage with such prior art dressings, however, is the fact that despite their smooth surface, they still get stuck in the wound bed much too often.

With the present invention there has been achieved a wound dressing which does not stick to healing wound tissue and which substantially does not give off fibers or other components harmfully affecting the wound healing process.

A wound dressing according to the invention is primarily distinguished in that it comprises a soft and elastic hydrophobic gel lying in direct contact with the wound bed during use of the dressing and being built up on a reinforcement netting made of an elastic material, said gel being applied to seal around all constituents of the netting while leaving through-holes in the layer composed of gel and reinforcement.

To accomplish a wound dressing having properties such as low tendency of sticking to wound tissue and poor solubility in aqueous media such as wound fluid for example, the gel incorporated in the dressing must necessarily be hydrophobic.

According to a particularly advantageous embodiment of the invention, the hydrophobic gel is a silicone gel.

The invention will be described in more detail below with reference to an exemplary embodiment illustrated in the accompanying drawing, of which

Fig. 1 is a plan view of a wound dressing according to the invention, and

Fig. 2 is a partially enlarged view of the dressing as shown in Fig. 1.

The dressing shown in Figs. 1 and 2 comprises a reinforcement 1 in the form of an elastic, hydrophobic, knitted network and a silicone gel layer 2 applied to seal around the threads of the reinforcement 1 which is thereby encapsulated by the gel. In order for the dressing to perform its optimal function, the encapsulation must be almost complete with only a strictly limited number of fiber ends being allowed to penetrate the silicone gel and project out of the dressing surface.

The gel must be securely fixed to the reinforcement 1 for maintaining the gel layer 2 substantially intact during use of the dressing, avoiding in this manner any disturbance of the wound healing process such as inflammatory reactions caused by loose gel fragments.

When, as in the example shown in Fig. 1, the gel seals around the structural elements included in the reinforcement 1 constituting here the individual threads 3 of the netting, it will form a continuous, three-dimensionally extended layer which is in itself strong enough to keep the reinforcement well encapsulated.

In order to provide a higher degree of security against disintegration of the gel layer, the reinforcing

threads 3 may, as shown in Fig. 2, be composed of a plurality of loosely twisted fibers 4. With such an arrangement the silicone gel will penetrate between each individual fiber in the threads of the netting, adding in this manner to the strength of the gel layer.

Thus, the gel does not necessarily have to adhere to the reinforcement material for the dressing to present a continuous and stable outer layer of gel.

The openings 5 in the net allows for the passage of excess wound exudate through the dressing. For the treatment of wounds under regeneration of skin tissue, a dressing according to the invention is preferably combined with an absorbent bandage applied over the gel-coated netting. In this manner excessive amounts of wound exudate will be sucked up at the same time as the absorbent bandage is kept spaced from the wound bed thereby preventing the bandage from sticking to the wound. If necessary, the distance between the absorbent bandage and the wound tissue can be further increased by superimposing several layers of gel-coated netting on the wound surface before applying the absorbent bandage.

Owing to its flexibility and its capability of adhering to dry skin, the silicone gel netting can be easily affixed over the wound site. The strongly hydrophobic nature of the silicone gel prevents wound fluid from penetrating out over the healthy skin surrounding the wound and to loosen up skin tissue. The net-like structure of the dressing allows for the surrounding, healthy area of the skin to which the dressing is affixed to be maintained airy and rich in oxygen.

The net coated with silicone is easily reshapable and self-adhering, allowing thereby several layers of the net to be applied in a superimposed position. Due to its pliability, the dressing can further

be used for filling in deep wound cavities enabling thereby the wound edges to be kept apart during the wound healing process. In this manner the wound is prevented from closing together by contraction of the wound edges while also reducing the risk of disfiguring and motion-inhibiting scars due to the gradual formation of new tissue from the wound edges, the wound cavity successively being filled in with regenerated tissue.

As an example of a silicone gel having the properties required for this purpose can be mentioned a type of gel marketed by Dow Corning under the registered tradename Dow Corning Q7-2218 - Silicone Gel System.

The hydrophobic character of the silicone gel makes it particularly useful as a carrier for medicaments soluble in fat such as pain-relieving substances, for example. It is also conceivable to have antibacterial agents, or agents stimulating wound repair, incorporated in the gel. An example of the last-mentioned type of agent is zinc.

The exemplary embodiment described in the foregoing is merely intended to illustrate the inventive concept without restricting its scope.

The silicone gel described in the example could naturally be replaced by any other hydrophobic gel having similar properties in other respects. For example, polyurethane gel could be used as an alternative to silicone gel.

Many modifications of the invention are conceivable within the scope of the patent claims.

In a suitable embodiment the reinforcement could be made of a polyurethane foam being immersed in a bath of silicone gel to accomplish the wound dressing.

CLAIMS

1. A wound dressing, characterized in that it comprises a soft and elastic hydrophobic gel (2) lying in direct contact with the wound during use of the dressing; that the hydrophobic gel is built up on a net-like reinforcement (1) made of an elastic material; and that the hydrophobic gel is applied to seal around all components of the network but leaving through-holes (5) in the layer built up of hydrophobic gel and reinforcement.
2. A wound dressing according to Claim 1, characterized in that the hydrophobic gel (2) is a silicone gel.
3. A wound dressing according to Claim 1, characterized in that the hydrophobic gel (2) is a polyurethane gel.
4. A wound dressing according to Claim 1, characterized in that the reinforcement (1) is a soft, flexible and elastically extendable net of textile material.
5. A wound dressing according to Claim 2, characterized in that the netting has a knitted, crocheted or woven performance.
6. A wound dressing according to any one of the preceding claims, characterized in that the netting has 5 - 30 holes per cm^2 , said holes (5) having a size of $0.25 - 4 \text{ mm}^2$.
7. A wound dressing according to Claim 1, characterized in that the reinforcement (1) consists of a polymeric foam with open cells.

1/1




FIG 1





FIG 2

INTERNATIONAL SEARCH REPORT

International Application No PCT/SE87/00113

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC :		
A 61 F 13/00, A 61 L 15/01		
II. FIELDS SEARCHED		
Minimum Documentation Searched :		
Classification System	Classification Symbols	
IPC 4	A 61 F 13/00, /02; A 61 L 15/00, /01, /06	
US C1	128: 82, 90, 155, 156, 390; 604: 304-310	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched *		
SE, NO, DK, FI classes as above		
III. DOCUMENTS CONSIDERED TO BE RELEVANT*		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	DE, A, 2 132 498 (S ROSOFF) 29 June 1972 see p 4 lines 3-8 & FR, 2100189	1-2, 4-5, 7
Y	FR, A, 2 285 112 (MOLINIER S A ET AL) 16 April 1976	1, 4-5
Y	US, A, 2 764 976 (W C SKILES ET AL) 2 October 1956	1-2, 4-5
A	US, A, 3 006 338 (T P DAVIES) 31 October 1961	1-2, 4-5
A	US, A, 3 113 568 (J S ROBINS) 10 December 1963	1, 3, 7
<p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
1987-05-25	1987-05-27	
International Searching Authority	Signature of Authorized Officer	
Swedish Patent Office	 Leif Karnsäter	